

































Jean-Louis Peyraud Deputy scientific director of Agriculture






# Part 1: A world without livestock is a nonsense



# A humanitarian, economic and agronomic nonsense

 Livestock provides livelihood to more than 800 million poor people

> Density of Poor Livestock Keepers Year 2010\*



- In smallholders family farming systems, livestock enhances food security
  - Nutritious food
  - organic fertilizer for crops & soils
  - workforce and transport (lack of roads)
  - source of regular income and savings
  - gender equity






# A nonsense for global food and nutrition security

 A lack of protein of animal origin cause anemia and stunting in some parts of the world

Food and Agriculture Organization of the United Nations



- Food from marginal lands?
   Ruminants can do!!!
  - In Europe, permanent grasslands and rangelands cover 73 M ha (40% Eu AA)





360 M cattle and 600 M small ruminants provide
 25% of world animal product from marginal land








# Livestock farming for a more efficient agriculture

#### We would feed more people without livestock: it's wrong!

- Complementarity between livestock and crops to maximize food production
  - Valorisation of co-products
  - Valorisation of uncultivated land



(Adapted from Van Kernebeck et al., 2014 et De Boeer et al., 2018)

More land is needed to feed a vegan population than a population eating 20 g of protein
of animal origin per day (1/3 of the current supply of animal protein)



Part 2:
Shadows and benefits
of livestock: beyond false
assumptions, towards a
more balanced vision



### Livestock between Food and Feed!


### "10 kg of plant proteins to produce 1 kg of animal proteins!"

Livestock consume 6 Billion tons dry matter, of which 86% are non edible as human food

Mottet et al., 2018

Kg of protein of animal origin per kg of edible plant protein used as feed


 Feed vs food competition does in fact concern those proteins of plant origin that are consumable by human (actually not consumed by animals)






### Real Carbon footprint of ruminants








### > 40% reduction

- Genotyping low methane production for selection
- Rumen microbiomes
- Improving animal health and husbandry conditions
- Smart use of manure
- More C sequestration (grassland, agroforestry)
- Precision Livestock Farming
- Feed production (circularity)
  - More efficient production (legumes)
  - Better agricultural land use (rotations)
  - Less/no specific feed production




# Meat consumption vs GHG emissions: where are the challenges?

Meat consumption



Emission intensity





CO.

## Is cow methane to blame for global warming?

Photochemical oxydation

Do not accumulate (10 years) so long emissions do not increase. Atmospheric concentration will decrease if emissions decrease

 $CH_4$ 

Photosynthesis
Soil Carbon

Using conventional GWPs to convert CH<sub>4</sub> to "CO<sub>2</sub>-eq" emissions misrepresents its impact on global temperature

Fossil CO<sub>2</sub> (100 years) accumulates in the atmosphere so long emissions exceed the ability of plants and ocean to take up new CO<sub>2</sub>






Fossil fuel
Old photosynthetic C
100-200 Million year old
(not in the C cycle)

- To limit warming to 1.5 to 2°C (COP 21)
  - CO<sub>2</sub> (and N<sub>2</sub>0) emissions should be reduce to zero,
  - CH<sub>4</sub> should be declining but do not have to reach net zero



# The ambivalent effect of livestock on biodiversity


 Intensive livestock contributes to biodiversity losses



- Ruminants produce biodiversity
  - Diversity of forage species (including honey plants) and grassland types



 Diversification of land uses, landscapes and maintenance of open habitats (with grasslands)









## Livestock provides ecosystem and social services

- Various benefits of a sustainable EU livestock sector for rural area
- The bundle of services varies according to local contexts
  - Intensive systems



Extensive systems






### **Designing animal-friendly systems**

#### Enrichment of the environment


#### % injured animals



Mirabito et Michel (2003)



Innovative buildings and precision livestock farming







# Part 3: Changing paradigms: towards a renewed place and role

of livestock farming in agri-food systems



#### The "linear" vision






- This as led to significant productivity gains but
  - In a linear way of thinking (resource -> production -> product -> waste)
  - Without considering the amount and origin of mobilized resources
  - Without preventing the degradation of ecosystems



### A new paradigm

#### Livestock is a key issue for sustainable circular agri-food systems



- Rethinking the place, roles and performances of livestock
- Rethinking the links between livestock, crop production, soil fertility and environment
- Rethinking the links between livestock, livestock products and consumption of animal based products
- Balances are to be found according to the political choices and the territorial contexts.


There is no « one size fits all » optimal solution



# Part 4: Take home messages




- Think twice: do not step into a simple and narrow vision of livestock farming systems
- Reducing impacts of livestock farming is essential:
   the shadow of livestock can be mitigated
- Livestock is not only a problem, it is also part of the solution for circular sustainable agri-food systems
- Livestock farming systems should change to regain legitimacy
- Europe needs an ambition for livestock farming systems: articulate local and global, transformation or improvement, food production and/or immaterial functions (multifunctional livestock)?






### **Local pollutions**



### Evolution of nitrate levels (mg / year) in groundwater (1998 – 2014)






### Water consumption by livestock

« 15 000 L of water per one kg of meat! »

- What are we talking about?
  - Green water (soil water consumed for crop cultivation): more than 95% is recycled
  - Blue water (surface water and groundwater)





- Livestock consume 8 to 15% of water resource worldwide (FAO, 2014)
- Comparison of farming systems

1 kg beef meat	22 – 520 L
1 kg pig/poultry meat	190 L
1 kg milk	< 1 - 100 L
1 shower	50 – 70 L

Doreau et al. (2014)